Index of Authors

with the titles of papers

ABE, H., *see* Makarov, V. I.
AGEEV, S. Z., and SANCTUARY, B. C.: Analytical solutions for spin 7/2 line intensities in solid state NMR, 835
AKINLADE, O., *see* Pastore, G.
ANDREWS, A. M., TRETYAKOV, M. YU., PATE, B. H., FRASER, G. T., and KLEINER, I.: Fourier-transform infrared and jet-cooled diode-laser spectra of the 867 cm^{-1} ν_9 band of acetaldehyde, 201
ANTA, J. A., LOMBA, E., MARTÍN, C., LOMBARDERO, M., and LADO, F.: A fast method of solving the hypernetted-chain equation for molecular Lennard-Jones fluids, 743
ANTA, J. A., and KAHL, G.: On the use of a non-additive reference system in a reference hypernetted chain calculation of the structure of a binary liquid, 1273

BADIRKHAN, Z., *see* Pastore, G.
BAGLIN, F. G., ROSE, E. J., and SWEITZER, S.: Identification of 1, 2 and 3 body Raman scattering by the field gradient induced dipole *A* tensor in methane, 115
BARKER, G. C., GRIMSON, M. J., and SILBERT, M.: Effective interactions in colloidal dispersions stabilized by free polymers, 211
BARTLETT, P., *see* Eldridge, M. D.
BAUDER, A., *see* Makarewicz, J.
BAUMANN, D., *see* Wu, J. Q.
BESNARD, M., *see* Tassaing, T.
BIBB, S. J., ROWLINSON, J. S., and XIAO, C.: Model of adsorption at a fluid–fluid interface, 1099
BORIN, A. C., ORNELLAS, F. R., and UENO, L. T.: The electric field gradient at the ^{9}Be nucleus in BeC, 133
BÖHM, M. C., *see* Schütt, J.
BRONDEAU, J., *see* Canet, D.
BROWN, D., and NEYERTZ, S.: A general pressure tensor calculation for molecular dynamics simulations, 577

CACCAMO, C., GIUNTA, G., and MALESCIO, G.: Microscopic theoretical determination of the phase diagram of the hard core Yukawa fluid, 125
CALLEJA, M., *see* Marsh, P.
CANET, D., MUTZENHARDT, P., and BRONDEAU, J.: A *p*-spin high-pass filter using radiofrequency field gradients for homonuclear magnetic resonance experiments, 597
CARTER, S., *see* Persson, B. J.
CHAKRABARTI, A., YASHONATH, S., and RAO, C. N. R.: A Monte Carlo study of the condensed phases of biphenyl, 49
CHAKRAVARTY, C.: Fourier path integral simulations of *para*-H_2 and *ortho*-D_2 clusters, 845
CHAMPENEY, D. C., *see* Okan, S. E.
CHIALVO, A. A., and CUMMINGS, P. T.: Comment on 'Near critical phase behaviour of dilute mixtures', 41
CIOSLOWSKI, J., and STEFANOV, B. B.: Variational determination of the zero-flux surfaces of atoms in molecules, 707
COSTA, C., *see* Figari, G.
CRIADO, A., *see* Muñoz, A.
CRUPI, V., JANNELLI, M. P., MAGAZU', S., MAISANO, G., MAJOLINO, D., MIGLIARDO, P., and SIRNA, D.: Rayleigh wing and Fourier transform infrared studies of intermolecular and intramolecular hydrogen bonds in liquid ethylene glycol, 645
CUMMINGS, P. T., *see* Chialvo, A. A.

0026–8976/95 $10·00 © 1995 Taylor & Francis Ltd.

DANTEN, Y., *see* Tassaing, T.
DE LORENZI, A., *see* Stoppa, P.
DE SANTIS, A., GREGORI, A., and ROCCA, D.: The corresponding states principle for radial distribution functions of homonuclear diatomic liquids, 1
DI BARI, L., MÄLER, L., and KOWALEWSKI, J.: Nuclear spin relaxation in chloroform dissolved in a nematic phase: spectral densities at high frequencies, 31
DELEEUW, B. J., YAMAGUCHI, Y., and SCHAEFER III, H. F.: Chromium dihydride (CrH_2): theoretical evidence for a bent $^5B_2$ ground state, 1109
DICKINSON, A. S., *see* Vesovic, V.
DING, F., *see* Fowler, P. W.
DORE, P., FINZI, L., NUCARA, A., POSTORINO, P., and ROVERE, M.: Translational absorption band in low density mixtures of noble gases: the He–Xe case, 1065
DUNMUR, D. A., *see* Osipov, M. A.

ELDRIDGE, M. D., MADDEN, P. A., PUSEY, P. N., and BARTLETT, P.: Binary hard-sphere mixtures: a comparison between computer simulation and experiment, 395
ELLIOTT, S. R., *see* Sergi, A.
ERNST, R. R., *see* Robyr, P.

FEDCHENIA, I. I., and WESTLUND, P.-O.: Energy transfer between two mobile chromophores in a zero energy transfer configuration. Cross-correlation effects in the fluorescence anisotropy $r(t)$, 159
FERRARIO, M., *see* Sergi, A.
FIGARI, G., SICILIANO, A., COSTA, C., and MAGNASCO, V.: On the convergence rate of a cusp-corrected damped multipole expansion for the second-order $H–H^+$ induction energy, 633
FIGARI, G., *see* Magnasco, V.
FINZI, L., *see* Dore, P.
FISCHER, J., *see* Kriebel, C.
FLORIÁN, J., LESZCZYŃSKI, J., and SCHEINER, S.: *Ab initio* study of the structure of guanine–cytosine base pair conformers in gas phase and polar solvents, 469
FOWLER, P. W., DING, F., and MUNN, R. W.: Polarizabilities of anions in anisotropic environments. The fluoride ion in the perovskite lattices $NaMgF_3$, $KMgF_3$ and $KCaF_3$, 787
FRASER, G. T., *see* Andrews, A. M.

GAMBI, A., *see* Stoppa, P.
GAZZILLO, D.: Stability of fluids with more than two components. II. Demixing ternary mixtures of non-additive hard spheres, 303
GHERSETTI, S., *see* Stoppa, P.
GIANTURCO, F. A.: Quantum calculations of transport and relaxation properties for the helium–methane system, 481
GIORGIANNI, S., *see* Stoppa, P.
GIUNTA, G., *see* Caccamo, C.
GÓŹDŹ, W. T., GUBBINS, K. E., and PANAGIOTOPOULOS, A. Z.: Liquid–liquid phase transitions in pores, 825
GREGORI, A., *see* de Santis, A.
GREINER, S. P., *see* Kim, B.
GRIMSON, M. J., *see* Barker, G. C.
GROB-PISANO, C., *see* Robyr, P.
GUBBINS, K. E., *see* Góźdź, W. T.
GUBBINS, K. E., *see* Ulberg, D. E.
GUILLOT, B., *see* Tassaing, T.
GUISSANI, Y., *see* Tassaing, T.

HAYASHI, H., *see* Igarashi, M.
HAYASHI, H., *see* Makarov, V. I.
HAYASHI, M., *see* Lin, S. H.
HAYMET, A. D. T., *see* Wang, J.

HEYES, D. M., and MITCHELL, P. J.: Non-equilibrium molecular and Brownian dynamics simulations of shear thinning of inverse power fluids, 261
HILLS, B. P., and SNAAR, J. E. M.: Water proton relaxation studies of pore microstructure in monodisperse glass bead beds, 141
HOLOVKO, M. F., and VAKARIN, E. V.: An associative version of the Henderson–Abraham–Barker equation, 1057
HORSEWILL, A. J., IKRAM, A., and TOMSAH, I. B. I.: Hydrogen bond dynamics in tetrafluoroterephthalic acid studied by NMR and INS, 1257
HUTSON, J. M., *see* Wheatley, R. J.
HUTSON, J. M.: Calculating nuclear quadrupole coupling constants for van der Waals complexes, 185

IGARASHI, M., MENG, Q.-X., SAKAGUCHI, Y., and HAYASHI, H.: Reversion of magnetic field effects under large magnetic fields observed in the photochemical hydrogen abstraction reactions of fluorinated acetophenone and benzophenone in micellar solutions, 943
IGEL-MANN, G., SCHLUNK, R., and STOLL, H.: Structure and ionization potentials of clusters containing heavy elements. Part 4: Mixed clusters of alkali and group VI elements, 679
IGEL-MANN, G., and STOLL, H.: Structure and ionization potentials of clusters containing heavy elements. Part 3: Mixed clusters of alkali and group V elements, 663
IKRAM, A., *see* Horsewill, A. J.

JANNELLI, M. P., *see* Crupi, V.
JAQUET, R., and RÖHSE, R.: Rovibrational energy levels and transitions for H_3^+ computed from a new highly accurate potential energy surface, 291
JEDLOVSZKY, P., and PÁLINKÁS, G.: Monte Carlo simulation of liquid acetone with a polarizable molecular model, 217

KAHL, G., *see* Anta, J. A.
KAMINSKY, R. D.: *N*-Particle configurational integrals of model fluid–micropore systems, 69
KAPLAN, I. G., SANTAMARIA, R., and NOVARO, O.: Non-additive forces in atomic clusters. The case of Ag_n, 105
KIM, B., GREINER, S. P., and KRELICK, R. W.: Magnetic angle selection in rhombic metal complexes and nitroxide radicals. EPR spectra as three-dimensional energy surfaces, 511
KITAO, O., *see* Yamamoto, R.
KLEINER, I., *see* Andrews, A. M.
KOCH, W., *see* Natterer, J.
KOLAFA, J., and NEZBEDA, I.: The hard tetrahedron fluid: a model for the structure of water?, 421
KOMOLKIN, A. V., and MALINIAK, A.: Local structure in anisotropic systems determined by molecular dynamics simulation. Application to a nematic liquid crystal, 1227
KOROCIŃSKI, A., and NAPIÓRKOWSKI, M.: Capillary interface potential and interfacial fluctuations, 171
KOWALEWSKI, J., *see* di Bari, L.
KRELICK, R. W., *see* Kim, B.
KRIEBEL, C., MÜLLER, A., WINKELMANN, J., and FISCHER, J.: Vapour–liquid equilibria of two-centre Lennard-Jones fluids from the *NpT* plus test particle method, 381

LADO, F., *see* Anta, J. A.
LEE, T. J., and RACINE, S. C.: A coupled-cluster study of the molecular structure, vibrational spectrum and relative energies of the XCN and XNC (X = F, Cl) isomers, 717
LESZCZYŃSKI, J., *see* Florián, J.
LIN, C. H., *see* Lin, S. H.
LIN, S. H., HAYASHI, M., LIN, C. H., YU, J., VILLAEYS, A. A., and WU, G. Y. C.: Theoretical studies of IR–UV sum frequency generation applied to adsorbed molecules, 453
LOMBA, E., *see* Anta, J. A.
LOMBARDERO, M., *see* Anta, J. A.
LU, B. C.-Y., *see* Tang, Y.

MADDEN, P. A., *see* Eldridge, M. D.
MAGAZÙ, S., *see* Crupi, V.
MAGNASCO, V., FIGARI, G., SICILIANO, A., and RUI, M.: One-centre multipole calculation of second-order perturbation energies for H_2^+, 813
MAGNASCO, V., *see* Figari, G.
MAISANO, G., *see* Crupi, V.
MAJOLINO, D., *see* Crupi, V.
MAKAREWICZ, J., and BAUDER, A.: Rovibrational Hamiltonian for molecular complexes, 853
MAKAROV, V. I., ABE, H., and HAYASHI, H.: External magnetic field acceleration of radiationless processes in the Ã state of gaseous oxalyl fluoride, 911
MALESCIO, G., *see* Caccamo, C.
MALINIAK, A., *see* Komolkin, A. V.
MARSH, P., RICKAYZEN, G., and CALLEJA, M.: The structure of a hard sphere fluid restricted by permeable walls, 799
MARTÍN, C., *see* Anta, J. A.
MATTHEWS, F., *see* Pastore, G.
MATYUSHOV, D. V., and SCHMID, R.: Charge separation/recombination reactions in non-polar fluids: a molecular description, 533
MÄLER, L., *see* di Bari, L.
MCCOURT, F. R. W., *see* Vesovic, V.
MCDONALD, I. R., *see* Sergi, A.
MCDONALD, P. J., and PINTER-KRAINER, M.: High pressure NMR study of methyl group tunnelling in dimethyl sulphide, 1021
MEDEROS, L., *see* Rascón, C.
MEIER, B. H., *see* Robyr, P.
MENG, Q.-X., *see* Igarashi, M.
MEYER, H.: Counterpropagating molecular beam scattering of $NH_3 + nH_2$, 1155
MIGLIARDO, P., *see* Crupi, V.
MITCHELL, P. J., *see* Heyes, D. M.
MORAIS, V. M. F., and VARANDAS, A. J. C.: Quasiclassical trajectory study of the $Li + Cs_2$ reaction, 957
MORRISS, G. P., *see* Peric, M.
MRYGLOD, I. M., OMELYAN, I. P., and TOKARCHUK, M. V.: Generalized collective modes for the Lennard-Jones fluid, 235
MUNN, R. W., *see* Fowler, P. W.
MUÑOZ, A., and CRIADO, A.: A molecular dynamics interpretation of the inelastic neutron spectra in the plastic phase of hexachloroethane, 1207
MUTZENHARDT, P., *see* Canet, D.
MÜLLER, A., *see* Kriebel, C.

NAKANISHI, K., *see* Yamamoto, R.
NAPIÓRKOWSKI, M., *see* Korociński, A.
NATTERER, J., and KOCH, W.: Quantum chemical predictions of the electron affinities of carbon–hydrogen clusters $C_{2n}H$, the CH binding energies and the gas phase acidities of polyacetylenes $C_{2n}H_2$ for $n = 1$–3, 691
NAVASCUÉS, G., *see* Rascón, C.
NEYERTZ, S., *see* Brown, D.
NEZBEDA, I., *see* Kolafa, J.
NOVARO, O., *see* Kaplan, I. G.
NUCARA, A., *see* Dore, P.

OKAN, S. E., SALMON, P. S., CHAMPENEY, D. C., and PETRI, I.: The solvation of cations in hydrogen-bonded molecular solvents: a neutron diffraction study on the structure of Ni^{2+} solutions in ethylene glycol and in glycerol, 325
OMELYAN, I. P., *see* Mryglod, I. M.
ORLIKOWSKI, T., *see* Paul-Kwiek, E.
ORNELLAS, F. R., *see* Borin, A. C.
OSIPOV, M. A., PICKUP, B. T., and DUNMUR, D. A.: A new twist to molecular chirality: intrinsic chirality indices, 1193

PÁLINKÁS, G., *see* Jedlovszky, P.
PACKER, M. J., and PICKUP, B. T.: An analysis of the magnetizability tensor at CHF level, 1179
PADILLA, P., and VEGA, C.: Packing effects on the conformational equilibrium of alkanes, 435
PANAGIOTOPOULOS, A. Z., *see* Góźdź, W. T.
PASTORE, G., MATTHEWS, F., AKINLADE, O., and BADIRKHAN, Z.: Numerical solution of the optimized random phase approximation, 653
PATE, B. H., *see* Andrews, A. M.
PAUL-KWIEK, E., and ORLIKOWSKI, T.: Quantum calculations of multistructure cross-sections for the Ca(4s5p, $^3P_2 \leftarrow {}^1P_1$) + He collisional process, 971
PEDERSON, L. A., and YARKONY, D. R.: On the spin–orbit induced radiationless decay of the $b^3\Sigma^-$ state of BH, 611
PERIC, M., and MORRISS, G. P.: Free energy and entropy production in planar Couette flow far from equilibrium, 1049
PERSSON, B. J., ROOS, B. O., and CARTER, S.: A theoretical study of the rovibronic spectra of CNO, NCO and CON, 619
PETERS, G. H., and VELASCO, E.: Influence of surface and torsion potentials on the melting properties of a hexane monolayer on a graphite substrate. A molecular dynamics simulation study, 1039
PETRI, I., *see* Okan, S. E.
PICKUP, B. T., *see* Osipov, M. A.
PICKUP, B. T., *see* Packer, M. J.
PINTER-KRAINER, M., *see* McDonald, P. J.
POPELIER, P. L. A., and STONE, A. J.: Formulae for the first and second derivatives of anisotropic potentials with respect to geometrical parameters (Erratum), 811
POSTORINO, P., *see* Dore, P.
PUSEY, P. N., *see* Eldridge, M. D.

RACINE, S. C., *see* Lee, T. J.
RAGHAVAN, K.: Molecular dynamics simulation of a needle–sphere binary mixture, 345
RAO, C. N. R., *see* Chakrabarti, A.
RASCÓN, C., NAVASCUÉS, G., and MEDEROS, L.: New exact relations for the many-cavity distribution function of non-uniform hard-sphere systems, 373
RICKAYZEN, G., *see* Marsh, P.
ROBYR, P., TOMASELLI, M., STRAKA, J., GROB-PISANO, C., SUTER, U. W., MEIER, B. H., and ERNST, R. R.: RF-driven and proton-driven NMR polarization transfer for investigating local order. An application to solid polymers, 995
ROCCA, D., *see* de Santis, A.
RÖHSE, R., *see* Jaquet, R.
ROOS, B. O., *see* Persson, B. J.
ROSE, E. J., *see* Baglin, F. G.
ROVERE, M., *see* Dore, P.
ROWLINSON, J. S., *see* Bibb, S. J.
RUI, M., *see* Magnasco, V.
RUSSELL, A. J., and SPACKMAN, M. A.: Vibrational averaging of electrical properties. Development of a routine theoretical method for polyatomic molecules, 1239

SAKAGUCHI, Y., *see* Igarashi, M.
SALMON, P. S., *see* Okan, S. E.
SANCTUARY, B. C., *see* Ageev, S. Z.
SANTAMARIA, R., *see* Kaplan, I. G.
SCHAEFER III, H. F., *see* DeLeeuw, B. J.
SCHEINER, S., *see* Florián, J.
SCHLICK, T., *see* Zhang, G.
SCHLUNK, R., *see* Igel-Mann, G.
SCHMID, R., *see* Matyushov, D. V.
SCHULTE, J., *see* Schütt, J.
SCHÜTT, J., SCHULTE, J., BÖHM, M. C., and SOOS, Z. G.: Resolution of the sign problem in quantum Monte Carlo simulations of annulenes, 1127

SEREBRENNIKOV, YU, A., and STEINER, U. E.: Nuclear spin–rotation interaction and non-Abelian gauge potential, 627
SERGI, A., FERRARIO, M., ELLIOTT, S. R., and MCDONALD, I. R.: Molecular dynamics study of the plastic–crystalline phase transition of tetraphosphorus triselenide, 727
SICILIANO, A., *see* Figari, G.
SICILIANO, A., *see* Magnasco, V.
SILBERT, M., *see* Barker, G. C.
SIRNA, D., *see* Crupi, V.
SNAAR, J. E. M., *see* Hills, B. P.
SOOS, Z. G., *see* Schütt, J.
SPACKMAN, M. A., *see* Russell, A. J.
STEFANOV, B. B., *see* Cioslowski, J.
STEINER, U. E., *see* Serebrennikov, Yu, A.
STEINER, U. E., *see* Wu, J. Q.
STOLL, H., *see* Igel-Mann, G.
STONE, A. J., *see* Popelier, P. L. A.
STOPPA, P., GIORGIANNI, S., GAMBI, A., DE LORENZI, A., and GHERSETTI, S.: High resolution FTIR spectrum of vinyl fluoride near 9 μm: rovibrational analysis of the ν_7 band, 281
STRAKA, J., *see* Robyr, P.
SUTER, U. W., *see* Robyr, P.
SWEITZER, S., *see* Baglin, F. G.

TANG, Y., and LU, B. C.-Y.: Analytical solution of the Ornstein–Zernike equation for mixtures, 89
TASSAING, T., DANTEN, Y., BESNARD, M., ZOIDIS, E., YARWOOD, J., GUISSANI, Y., and GUILLOT, B.: A far infrared study of water diluted in hydrophobic solvents, 769
THACHUK, M., *see* Vesovic, V.
TOKARCHUK, M. V., *see* Mryglod, I. M.
TOMASELLI, M., *see* Robyr, P.
TOMSAH, I. B. I., *see* Horsewill, A. J.
TRETYAKOV, M. YU., *see* Andrews, A. M.

UENO, L. T., *see* Borin, A. C.
ULBERG, D. E., and GUBBINS, K. E.: Water adsorption in microporous graphitic carbons, 1139

VAKARIN, E. V., *see* Holovko, M. F.
VARANDAS, A. J. C., *see* Morais, V. M. F.
VEGA, C., *see* Padilla, P.
VELASCO, E., *see* Peters, G. H.
VESOVIC, V., WAKEHAM, W. A., DICKINSON, A. S., MCCOURT, F. R. W., and THACHUK, M.: Quantum mechanical calculation of generalized collision cross-sections for the He–N_2 interaction. Part II. Thermomagnetic effect, 553
VILLAEYS, A. A., *see* Lin, S. H.

WAKEHAM, W. A., *see* Vesovic, V.
WANG, J., and HAYMET, A. D. J.: Theory of an excess electron in fluid helium. The effect of non-pairwise additivity of the solvent-induced interactions, 1033
WESTLUND, P.-O., *see* Fedchenia, I. I.
WHEATLEY, R. J.: The non-additive exchange energies of H_3 and He_3, 899
WHEATLEY, R. J., and HUTSON, J. M.: A systematic model potential for Li^+–H_2O, 879
WINKELMANN, J., *see* Kriebel, C.
WU, G. Y. C., *see* Lin, S. H.
WU, J. Q., BAUMANN, D., and STEINER, U. E.: Interference of heavy-atom with magnetic spin effects in spin-correlated micellar radical pairs, 981

XIAO, C., *see* Bibb, S. J.

YAMAGUCHI, Y., *see* DeLeeuw, B. J.
YAMAMOTO, R., KITAO, O., and NAKANISHI, K.: Can the 'van der Waals loop' vanish? II. Effect of domain size, 757
YARKONY, D. R., *see* Pederson, L. A.
YARWOOD, J., *see* Tassaing, T.
YASHONATH, S., *see* Chakrabarti, A.
YU, J., *see* Lin, S. H.

ZHANG, G., and SCHLICK, T.: Implicit discretization schemes for Langevin dynamics, 1077
ZOIDIS, E., *see* Tassaing, T.